

Research



Cite this article: Mohring B, Angelier F, Jaatinen K, Steele B, Lönnberg E, Öst M. 2022 Drivers of within- and among-individual variation in risk-taking behaviour during reproduction in a long-lived bird. *Proc. R. Soc. B* **289**: 20221338.

<https://doi.org/10.1098/rspb.2022.1338>

Received: 10 July 2022

Accepted: 30 August 2022

Subject Category:

Behaviour

Subject Areas:

behaviour, ecology

Keywords:

common eider, *Somateria mollissima*, flight initiation distance, phenotypic plasticity, predation risk, selection hypothesis

Author for correspondence:

Bertille Mohring

e-mail: bertille.mohring@abo.fi

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.6182022>.

Drivers of within- and among-individual variation in risk-taking behaviour during reproduction in a long-lived bird

Bertille Mohring^{1,2}, Frédéric Angelier², Kim Jaatinen³, Ben Steele⁴, Elin Lönnberg⁵ and Markus Öst^{1,6}

¹Environmental and Marine Biology, Åbo Akademi University, 20500 Turku, Finland

²Centre d'Etudes Biologiques de Chizé, UMR 7372 CNRS – La Rochelle Université, 79360 Villiers-en-Bois, France

³Nature and Game Management Trust Finland, 10160 Degerby, Finland

⁴School of Arts and Sciences, Colby-Sawyer College, New London, NH 03257, USA

⁵S:t Olofsskolan, 20700 Turku, Finland

⁶Novia University of Applied Sciences, 10600 Ekenäs, Finland

BM, 0000-0001-8553-4041; FA, 0000-0003-2619-167X; KJ, 0000-0002-5568-5647; MÖ, 0000-0002-2205-1437

Plastic and selective mechanisms govern parental investment adjustments to predation threat. We investigated the relative importance of plasticity and selection in risk-taking propensity of incubating female common eiders *Somateria mollissima* facing unprecedented predation in SW Finland, Baltic Sea. Using a 12-year individual-based dataset, we examined within- and among-individual variation in flight initiation distance (FID), in relation to predation risk, nest detectability, individual traits and reproductive investment ($N_{\text{FID}} = 1009$; $N_{\text{individual}} = 559$). We expected females nesting in riskier environments (higher predation risk, lower nest concealment) to mitigate environmentally imposed risk by exhibiting longer FIDs, and females investing more in current reproduction (older, in better condition or laying larger clutches) to display shorter FIDs. The target of predation—adult or offspring—affected the mechanisms adapting risk-taking propensity; females plastically increased their FID under higher adult predation risk, while risk-avoiding breeders were predominant on islands with higher nest predation risk. Risk-taking females selected thicker nest cover, consistent with personality-matching habitat choice. Females plastically attenuated their anti-predator response (shorter FIDs) with advancing age, and females in better body condition were more risk-taking, a result explained by selection processes. Future research should consider predator type when investigating the fitness consequences of risk-taking strategies.

1. Background

Life-history theory predicts that prey should adjust their parental investment to the perceived predation threat [1]. Reproduction involves a trade-off between investing energy and resources in producing and caring for offspring, and saving resources for self-maintenance and survival [1–3]. This survival-reproduction trade-off is predicted to be stronger when parents are exposed to elevated predation risk [4]. To cope with predation threat, prey have developed anti-predator behaviours through natural selection and behavioural plasticity [5,6], both reflecting the fundamental trade-off between reproduction and survival.

Predator-prey interactions shape prey anti-predator traits and behaviours over time and space through selection processes [7,8]. Thus, lower quality or risk-taking prey may suffer higher predation mortality and/or reduced reproductive output under threat [9–11]. Such differences in survival or reproduction may induce phenotypic variation in survival rates and population

dynamics [12–15]. However, individuals may also plastically modulate their behavioural or physiological anti-predator responses depending on their internal state and extrinsic factors, including the level of nest and adult predation risk [7,16,17]. Consequently, phenotypic plasticity can confound patterns of selective (dis)appearance of risk-taking phenotypes, highlighting the need to account for both plasticity (within-individual processes) and selection (among-individual processes) in order to disentangle those responsible for population-level variation [18]. Identifying the drivers of variation in risk-taking behaviours is thus key to understanding their consequences for both life-history evolution and population dynamics, especially in a context of fluctuating predation threat. However, few studies have attempted to separate plasticity and selection of risk-taking phenotypes, given that such analyses require large longitudinal datasets covering a substantial part of the lifespan of the study organism, which are often difficult to obtain in the wild [19].

Flight initiation distance (FID), the distance at which prey start to flee from approaching potential predators [20], is a standardized measure to investigate risk-taking propensity and the relative roles of plasticity and selection in modulating risk-taking behaviour under fluctuating predation threat [21]. FID has been shown to be repeatable within individuals [22–26]. On the one hand, fleeing from the nest may jeopardize reproductive success by disrupting incubation and by indicating offspring location to the approaching nest or adult predator [27]. On the other hand, delaying escape may reduce breeder survival [28]. Consequently, individuals investing more in the current reproductive event are expected to dampen their anti-predator response and delay escape to enhance offspring survival [29,30].

The evolutionary consequences of individual variation in risk-taking under predation threat depend on the relative magnitude of adaptive phenotypic plasticity (i.e. phenotypes actively matching prevailing risk) and selective disappearance effects (i.e. exclusion of certain phenotypes from the breeding pool) [31–33]. Our long-term longitudinal data on FIDs of incubating female common eiders *Somateria mollissima*, hereafter eiders, allowed us to improve knowledge on the relative influence of these mechanisms on risk-taking behaviour during reproduction. Specifically, we investigated how FID is modulated by phenotypic traits, parental investment, conspicuousness to predators and fluctuating predation pressure. The study was conducted amidst an unprecedented increase in predation pressure in our study population in the northern Baltic Sea, mainly attributed to the recent recovery of the white-tailed eagle *Haliaeetus albicilla*, the main predator of adult eiders [34]. This created an ideal backdrop to investigate individual plasticity and population-level selection processes involved in prey responses to a fluctuating predation regime [35].

To this end, we first delineated the population-level trends in FID, female and nest characteristics, clutch size and predation risk over the entire study period (2008–2019). We then addressed whether FID was (i) related to female phenotypic traits, reproductive investment, nest conspicuousness and predation risk, (ii) repeatable and (iii) shaped by within-individual plasticity or among-individual selection. We assumed that FID would be affected by population-level selection against risk-taking phenotypes (i.e. displaying shorter FIDs) under the progressively increasing predation threat. In addition and according to life-history theory [1],

we expected females to be more risk-taking, i.e. display shorter FIDs, with advancing age (diminishing residual reproductive value), when the costs of reproduction are lower (good body condition) and when making a greater reproductive investment (large clutch). Given that nest conspicuousness can affect predation risk, we expected females to advance their escape response when breeding in nests offering little concealment, or on islands characterized by high adult and nest predation risk. Alternatively, but not exclusively, females characterized by shorter FIDs may avoid conspicuous nests and risky islands, hence mitigating environmentally imposed risk of predation. We further expected FID to be individually repeatable, but modulated by plastic adjustments to individual state, parental investment and perceived predation risk.

2. Methods

(a) Female monitoring

(i) Breeding female characteristics

The study was conducted in the western Gulf of Finland, in the archipelago surrounding Tvärminne Zoological Station (59°50'N, 23°15'E), in 2008–2019. Female eiders nest on diverse types of islands, including those covered by bare rocks, sparse patches of herbs, stands of juniper *Juniperus communis* or pine *Pinus sylvestris* dominated forest. Incubating female eiders at Tvärminne have been annually trapped on their nest with hand nets since 1990, and individually marked with a standard metal ring. Female weight was measured to the nearest 10 g with a Pesola spring balance, and radius-ulna length, a proxy of body size, was measured to the nearest 1 mm with a wing ruler. Females incubate for about 26 days [36], and females were trapped between May and early June during the later phase of incubation to minimize nest abandonment [37]. Hatching date was estimated using an egg floatation test [38], and the number of eggs was counted as a proxy of initial energetic investment in reproduction [39,40].

Eiders are long-lived (*ca* 21 years [41]) and philopatric [42,43] birds periodically relying on intermittent breeding [44]. Females were ringed when caught breeding for the first time, which typically occurs at three years-of-age [45]. As trapping success is high and trapping effort has been constant since 1996 [46], we used the number of years since the bird was first trapped and ringed as a proxy of female minimum age, also reflecting residual reproductive value. As capital breeders, eiders fast and rely on their body reserves both to produce eggs and for maintenance during incubation [47]. Maternal energy reserves are progressively depleted as incubation proceeds, necessitating the need to assess the amount of body reserves at a comparable time. We therefore estimated a body condition index at hatching, which corresponds to the time of FID measurement (see electronic supplementary material S1, [48]).

(ii) Risk-taking propensity

Animals usually respond to human approach as they would to an approaching natural predator [49]. We thus measured risk-taking propensity with a standardized FID protocol consisting of a unique observer approaching the nest at a constant slow gait and measuring the distance at which the incubating focal female fled [50]. Before starting the approach, the observer positioned himself so that the female had a free line of sight to him, ensuring that the focal female had noticed him and followed him with her gaze or slight head movements. Because a pilot study in 2012 showed that starting distance was not correlated with FID in this population (see electronic supplementary material S2),

Table 1. Number of FID measurements per incubating female common eider.

number of FID replicates (2008–2019)	1	2	3	4	5	6	7	8	9
number of females	310	131	73	23	13	5	2	1	1

starting distance was not recorded. As FID often decreases when the clutch nears hatching [28,51,52] but also to limit the risk of nest abandonment, FID was measured the day before the expected hatching date predicted by egg floatation. Once the female had fled from the nest, the observer immediately covered the nest to minimize the risk of detection and depredation by avian egg predators such as hooded crows *Corvus cornix* [53]. On average, $33.08 \pm 5.94\%$ of nests active at first encounter were annually sampled for FID (range: 24.19%–42.92%). This relatively low proportion was due to the fact that females whose eggs were depredated or hatched prior to FID measurement could not be sampled for FID ($61.74 \pm 13.79\%$ of successful nests were on average annually sampled, range: 43.70%–87.50%). Coupled with a predation risk-induced increase in intermittent breeding strategies and premature mortality due to predation [34], the number of repeated measures per individual over time was lower than what one could expect from such a long-lived philopatric species (table 1). Overall, we measured 1009 FIDs to the closest 10 cm (later rounded to the closest meter (see §2(c)) to account for measurement imprecision and to meet model assumptions) on 559 incubating females over the 12-year study period.

(b) Perceived predation risk

(i) Nest concealment

Nest concealment was determined for each nest based on hemispherical photographs, aimed from the centre of each nest. Photographs were taken using a 42 mm semi-fisheye lens mounted on an Olympus C-740 digital camera [48], covering lateral and vertical nest surroundings. Each picture was converted to black and white, and the proportion of black pixels (corresponding to rocks or vegetation concealing the nest) was calculated using ImageJ [54].

(ii) Island-specific adult and nest predation risk

The main local predators of breeding females are the white-tailed eagle, the American mink *Neovison vison* and the raccoon dog *Nyctereutes procyonoides* [34,55]. In addition to predation on breeding females, eider eggs can be consumed either by primary predators of female eiders, or by avian predators like hooded crows or gulls *Larus* spp., usually depredating eggs during female absence from the nest [56]. In the study area, the numbers of killed females and depredated nests vary greatly over time and space (see electronic supplementary material S3). We thus calculated yearly indices of adult and nest predation risk for each of the 19 study islands over the 12-year study period.

First, island-specific adult predation risk index was defined as the proportion of killed nesting females, obtained by dividing the number of killed females on a given island and year by the number of breeding attempts on this island and year (see [55]). Breeding attempts included active nests ($N = 4097$ in 2008–2019), nests that had already been depredated ($N = 878$), and nests in which ducklings had already hatched ($N = 172$).

Second, island-specific nest predation risk index was obtained by dividing the number of failed breeding attempts at the final nest fate census on a given year and island by the number of nests on the island whose fate (success or failure) was known [57]. To this aim, we monitored each nest fate (failure or success) at the end of the breeding period. Eiders are precocial

birds and ducklings usually leave the nest within a day after hatching [58]. Successful nests were characterized by the presence of ducklings or hatched eggshells, recognizable by a leathery membrane. Depredated nests were characterized by the absence of eggshells or the presence of shells broken into pieces with a thin bloody membrane [48]. Nest failure was mainly due to depredation, and non-depredated but failed nests could be explained by females abandoning their nest or, occasionally, by females incubating unviable eggs. In some cases, nest depredation may result from the nest being depredated after the female had been killed, but as the number of depredated nests exceeded by many folds the number of females found killed and as the majority of females are observed in the study area after breeding failure, we believe that female depredation is not the main explanation for nest depredation. In addition, the correlation between our two island-specific predation risk indices was relatively low and variable over the study period (mean yearly correlation \pm s.d. = 0.30 ± 0.19 , range 0.05–0.62, see electronic supplementary material S4), suggesting that the two processes may be relatively independent from each other.

(c) Statistical analyses

Statistical analyses were conducted in R v.4.0.2. [59]. For all models, we ensured that the assumptions of normality and homoscedasticity of residuals were met. FID values were rounded to the closest meter to meet model assumptions and all explanatory variables were centred and scaled. We also checked for overdispersion after fitting Poisson generalized linear mixed models (GLMMs) [60]. We did not detect multi-collinearity among explanatory variables in any model (variance inflation factors, $VIFs < 2.5$ [61]).

First, we implemented GLMMs and linear mixed models (LMMs) to assess population-level trends in focal variables ($N_{FID} = 1009$; $N_{individual} = 559$; $N_{island} = 19$). Female and island identity were included as random effects to account for the non-independence of observations from the same individual (FID, female minimum age, body condition, nest cover and clutch size) and/or island (all focal variables). GLMMs with a Poisson distribution were used to study annual variation in FID and female minimum age, and LMMs to assess variation in body condition, nest cover, clutch size, and island-specific nest and adult predation risk.

Second, we constructed a GLMM to investigate variation in FID in relation to environmental variables and intrinsic individual characteristics ($N_{FID} = 1009$; $N_{individual} = 559$). We constructed a Poisson GLMM with FID as the dependent variable and female age, body condition, nest cover, clutch size and island-specific adult and nest predation risk as explanatory variables. Female and island identity were included as random effects to account for the non-independence of observations from the same individual and island.

Third, to elucidate the relative contributions of within- and among-individual changes in the explanatory variables on FID, we used within-subject centring [18,62] on a subset of individuals measured at least twice ($N_{FID} = 691$; $N_{individual} = 245$). Each explanatory variable was partitioned into its within-individual component, calculated by mean-centring the focal variable within each individual, and its among-individual component, calculated as the mean value of the focal variable for each




Figure 1. Yearly fluctuations in female eider (a) flight initiation distance, (b) clutch size, (c) minimum age, (d) body condition index, (e) nest cover and island-specific (f) nest and (g) adult predation risk over the 12-year study period. Black dots correspond to yearly population-level mean values of the focal variables and bars account for yearly population-level standard errors.

observed individual (see equation 2 in [62]). The within-individual component describes how each repeated observation deviates from the individual's mean (suggesting plastic behavioural responses) and the among-individual component describes how each individual differs from the population mean (suggesting non-random distribution of phenotypes potentially induced by directional selection on the focal trait) [62]. We thus constructed a Poisson GLMM with FID as the dependent variable and within-individual and among-individual components of age, body condition, nest cover, clutch size, and island-specific adult and nest predation risk as explanatory variables, and female and island identity as random effects. We further calculated adjusted within-individual repeatability of FID from this model [63]. Additionally, when both within- and among-individual effects were significant, we tested whether the within- and among-individual effects significantly differed from each other (see equation 3 in [62]). Within-individual

variance and repeatability of explanatory variables are further detailed in the electronic supplementary material S5 and S6.

3. Results

(a) Population-level temporal trends in focal variables

Annual fluctuation in the focal variables is presented in figure 1, and further description of the variable-specific variation over time is given in the electronic supplementary material S7. FID decreased over time (GLMM: estimate \pm standard error (E \pm s.e.) = -0.245 ± 0.034 ; $z = -7.169$, $p < 0.001$, electronic supplementary material S7A), as did clutch size (LMM: E \pm s.e. = -0.096 ± 0.039 , $t = -2.436$, $p = 0.015$, electronic supplementary material S7B). On the contrary, female minimum age

(GLMM: $E \pm \text{s.e.} = 0.586 \pm 0.032$, $z = 18.246$, $p < 0.001$, electronic supplementary material S7C), body condition (LMM: $E \pm \text{s.e.} = 0.155 \pm 0.028$, $t = 5.604$, $p < 0.001$, electronic supplementary material S7D) and nest cover (LMM: $E \pm \text{s.e.} = 0.036 \pm 0.006$, $t = 5.719$, $p < 0.001$, electronic supplementary material S7E) showed a positive time trend, as did nest (LMM: 0.067 ± 0.016 ; $t = 4.080$, $p < 0.001$, electronic supplementary material S7F) and adult (LMM: 0.024 ± 0.008 ; $t = 3.074$, $p = 0.002$, electronic supplementary material S7G) predation risk.

(b) Intrinsic and extrinsic determinants of flight initiation distance

Shorter FIDs were associated with advancing female (minimum) age (table 2, figure 2*a*), better body condition (table 2, figure 2*b*) and greater nest concealment (table 2, figure 2*c*). By contrast, increased island-specific adult predation risk was linked to longer FIDs (table 2, figure 2*d*).

(c) Flight initiation distance repeatability and within- and among-individual variation in flight initiation distance predictors

FID was significantly and moderately repeatable within individuals (adjusted repeatability, $r = 0.400$, 95% confidence interval, CI = [0.317, 0.527]). Shorter FIDs were associated with advancing age (table 3). As the within-individual slope was not steeper than the between-individual slope (non-significant difference between among- and within-individual slopes; $z = 1.121$; $p = 0.262$; see equation 3 in [62]), our results are in favour of a plastic response to ageing, but do not allow us to confirm selective disappearance of old breeders with long FIDs [64]. Shorter FIDs were also associated with good body condition and high nest concealment, these effects occurring only on the among-individual level (table 3). Interestingly, although high island-specific adult and nest predation risk were both significantly associated with longer FIDs, the FID response to adult predation risk was due to within-individual variation (table 3), whereas that to nest predation risk was confined to the among-individual level (table 3).

4. Discussion

Here, we investigated the patterns and drivers of variation in a standardized measure of risk-taking propensity (FID) in incubating female eiders experiencing increasing predator-induced mortality over 12 years. The temporal increase in adult and nest predation risk was accompanied by an increase in nest concealment, which presumably reflects the combined effect of adaptive responses, such as behavioural compensation to reduce detection by the visually hunting white-tailed eagle, and direct predator-induced selection against open nesting (see §4(a)). Contrary to our expectations, a concomitant decrease, rather than increase, in FID was observed, conceivably due to the action of overcompensatory intrinsic and extrinsic mechanisms (see §4(a)). Our most novel finding was that the prime target of predation—adult versus offspring—had a profound influence on individual variation in escape decision. Indeed, females plastically adjusted their risk-taking propensity to the level of perceived threat to themselves, while individuals with different risk-taking profiles settled on islands characterized by different

Table 2. GLMM explaining variation in FID in relation to individual (minimum age, body condition, nest cover and clutch size) and environmental (island-specific adult and nest predation risk) explanatory variables. Female and island identity were included as random effects. Significant parameter estimates ($p < 0.05$) are in italics.

fixed effect	estimate \pm s.e.	<i>z</i>	<i>p</i>
intercept	0.63 \pm 0.09	6.713	<0.001
<i>female minimum age</i>	<i>-0.46 \pm 0.04</i>	<i>-11.736</i>	<i><0.001</i>
<i>body condition index</i>	<i>-0.08 \pm 0.03</i>	<i>-2.599</i>	<i>0.009</i>
<i>nest cover</i>	<i>-0.13 \pm 0.04</i>	<i>-3.220</i>	<i>0.001</i>
clutch size	-0.00 \pm 0.3	-0.069	0.943
<i>island-specific adult predation risk</i>	<i>0.07 \pm 0.03</i>	<i>2.104</i>	<i>0.035</i>
island-specific nest predation risk	-0.00 \pm 0.3	-0.071	0.945

levels of threat to their nest. Thus, risk-avoiding phenotypes were predominant on islands with high nest predation risk, while breeding females plastically adjusted their anti-predator behaviour to perceived predation threat on themselves. These differential predator type-specific mechanisms linking FID responses to predation threat are consistent with the fundamental life-history trade-off between reproduction and survival (see §4(c)).

(a) Population-level temporal trends in focal variables

Intriguingly, and in contrast with findings in other species [65–67], higher adult and nest predation risk correlated with lower—rather than higher—population-level FID. The statistically significant decrease in FID over the study period is not visible when viewing mean annual population-level FID (figure 1*a*), becoming evident only by the use of random effects, which correct for pseudoreplication and provide a phenotypic estimate that blends within- and among-individual effects [68]. Two co-occurring compensatory mechanisms could mask and even negate the detection of a population-level increase in FID in response to increasing threat. First, nest cover has significantly increased over the study period, which, all else equal, should translate into shorter FIDs (figure 2*c*). This temporal shift in nest concealment likely represents an adaptive response to escalating predation by the visually hunting white-tailed eagle, which prefers open habitats, in which incubating females consequently suffer higher mortality [34,69]. At the macrohabitat scale, nesting on open islands may gradually decline because of higher predation [69,70] and lower breeding propensity linked to increased predation risk [34,71]. However, because females are highly philopatric to their nesting site [42,43], also behavioural compensation through microhabitat selection of concealed nest-sites within islands is likely involved. Second, consistent with previous findings, female body condition has increased over time [34,55] and in addition, the population has aged [55], probably contributing to a population-level shift towards shorter FIDs (figure 2*a,b*). The temporal increase in body condition of breeders may partly be due to more frequent predation-induced nest failure




Figure 2. Relationship between predicted flight initiation distance of incubating female eiders and (a) female age, (b) body condition, (c) nest concealment and (d) island-specific adult predation risk. Black dots correspond to mean values of the explanatory variable over regular intervals with dot size being proportional to the number of available data points for each interval, and bars account for standard errors. Regression lines depict the population response without integrating random effects, while the grey areas account for 95% confidence intervals.

by poor-condition individuals before monitoring onset, as good body condition is associated with higher nest success [57,72–74]. Poor-condition individuals may also be more likely to refrain from breeding [34,72] or suffer higher predation-induced mortality [69]. For example, in blue petrels *Halobaena caerulea*, poor body condition early in the season was associated with lower breeding success and a higher proportion of non-breeders [72]. In addition, increasing nest predation risk may reduce the production and recruitment of new breeders, and eventually translate into the observed ageing of the population (electronic supplementary material S7C), which has also previously been identified [55].

(b) Parental investment and phenotypic traits as drivers of plasticity and selection in risk-taking behaviour

The moderate within-individual repeatability of FID ($r = 0.400$) leaves room for differential selective disappearance of individuals depending on risk-taking propensity, but also for individually flexible adjustments of anti-predator behaviours according to internal state and environmental

conditions. The repeatability of FID is at the lower end of published estimates, typically ranging from 0.34 to 0.88 [22–26], and lower than the one previously reported for eiders over a shorter portion of the species' lifespan [50], probably reflecting the fact that repeatability tends to decline with time between samplings [75].

Two mutually non-exclusive mechanisms could explain the dampened anti-predator response with increasing individual quality and parental investment (age and body condition effects, table 2, figure 2). First, empirical evidence suggests that individuals in better health or condition tolerate closer approaches from potential predators [76,77]. Thus, assuming that shorter FIDs are correlated with higher individual quality, selection processes could explain the among-individual correlation between good body condition and short FIDs (figure 2b). Higher-quality individuals may face a lower rate of predation (e.g. due to possessing superior escape abilities [78,79]). Indeed, the survival of female eiders increases with body condition in our study population [69]. Selective disappearance effects may also be manifested through differences in breeding propensity. Increased predation risk is linked to

Table 3. GLMM explaining variation in FID in relation to within- and among-individual changes in explanatory variables (minimum age, body condition, nest cover, clutch size and island-specific adult and nest predation risk). Female and island identity were included as random effects. Significant parameter estimates ($p < 0.05$) are in italics.

fixed effect	estimate \pm s.e.	z	p
intercept	0.49 \pm 0.12	4.266	<0.001
<i>age_{Within}</i>	<i>-0.26 \pm 0.03</i>	<i>-9.062</i>	<i><0.001</i>
<i>age_{Among}</i>	<i>-0.34 \pm 0.07</i>	<i>-4.810</i>	<i><0.001</i>
body condition _{Within}	-0.04 \pm 0.03	-1.340	0.180
<i>body condition_{Among}</i>	<i>-0.13 \pm 0.06</i>	<i>-2.185</i>	<i>0.029</i>
nest cover _{Within}	-0.03 \pm 0.02	-1.277	0.202
<i>nest cover_{Among}</i>	<i>-0.25 \pm 0.07</i>	<i>-3.306</i>	<i><0.001</i>
clutch size _{Within}	0.03 \pm 0.03	1.233	0.217
<i>clutch size_{Among}</i>	<i>-0.01 \pm 0.06</i>	<i>-0.247</i>	<i>0.805</i>
<i>island-specific adult</i>	<i>0.10 \pm 0.03</i>	<i>3.445</i>	<i><0.001</i>
<i>predation risk_{Within}</i>	<i>0.02 \pm 0.07</i>	<i>0.288</i>	<i>0.77</i>
<i>predation risk_{Among}</i>	<i>-0.02 \pm 0.03</i>	<i>-0.641</i>	<i>0.521</i>
<i>island-specific nest</i>	<i>0.15 \pm 0.03</i>	<i>2.404</i>	<i>0.016</i>
<i>predation risk_{Among}</i>	<i>0.15 \pm 0.03</i>	<i>2.404</i>	<i>0.016</i>

a higher incidence of non-breeding in long-lived species [34,70,71], and we may hypothesize that risk-avoiding breeders with low body reserves (associated with longer FIDs), are particularly reluctant to jeopardize their own survival prospects and therefore more frequently skip breeding. A similar effect was not found for females laying larger clutches, but the low repeatability of clutch size ($r = 0.072$; electronic supplementary material S6) may not allow us to detect a potential among-individual effect of clutch size on FID.

Second, females could plastically adjust their risk-taking propensity to their level of parental investment. Females should be more willing to take risks to favour current reproduction as they age and the prospects of future reproduction diminish ('restraint hypothesis' [80]), supported by ample empirical evidence [7,26,29,81]. Indeed, females plastically decreased their FID with advancing age (figure 2a). However, we also expected current brood value to be higher for good-condition mothers and for females laying larger clutches, reflected in a reduced FID response. This argument hinges on the fact that mothers in good body condition produce larger clutches [82,83], incubate more constantly [84,85] and guard offspring more intensely [86]. Intriguingly, we did not find any evidence of females plastically adjusting their FID to their current body condition or clutch size. One plausible explanation could be that body condition is not as individually labile as often envisaged [73,86]. Supporting this hypothesis, female body condition was repeatable within individuals (repeatability, $r = 0.484$; electronic supplementary material S6). Similarly, undetected partial clutch predation, potentially contributing to the temporal association between decreasing clutch size and increasing nest predation risk, may obscure

the link between actual reproductive investment and maternal anti-predator responses. Conspecific brood parasitism may also be involved [87,88], although it is only a marginal phenomenon in this low-density population [87].

(c) Predation as a driver of plasticity and selection in risk-taking propensity

Females breeding in concealed nests displayed shorter FIDs. This confirms previous evidence that well-hidden prey in sheltered habitats tolerate a closer approach before fleeing [28,52,89–92], including eiders [93]. However, there is ongoing debate on whether individuals adjust their behaviour according to their local environment [94], or settle in habitats providing the best match to their behavioural profile ('personality-matching hypothesis' [95]), or phenotype in general [32]. Our results are more consistent with the personality-matching hypothesis. We conclude this because female eiders did not plastically adjust their anti-predator behaviour to their level of nest concealment over years (no within-individual effect), while there was a predominance of risk-taking females in concealed nests (significant among-individual effect). Our findings parallel those of D'Alba *et al.* [96]: female eiders varying in their physiological stress-coping phenotypes preferred nests with different shelter, rather than nest shelter directly affecting maternal stress levels. However, because the low within-individual variation in nest cover (electronic supplementary material, table S2) may prevent the detection of a significant within-individual effect of variation in nest cover on FID, our findings of support for the personality-matching hypothesis should be interpreted with some caution.

We provide, to our knowledge, the first demonstration that the target of predation determines the mechanisms adapting risk-taking propensity to the perceived level of threat. Thus, females plastically adjusted their escape responses to the perceived predation risk on themselves but not on their nests, whereas nest predation risk affected the relative distribution of risk-taking phenotypes in the population. Fundamental life-history trade-offs may explain these seemingly counterintuitive findings. Adult survival in long-lived iteroparous animals is typically high and canalized against environmental perturbations, and thus species should prioritize their own survival at the expense of current reproductive success (e.g. [97,98]). Higher individual risk should therefore advance escape responses, as indeed found here (figure 2d). However, the low average individual variance in island-specific nest predation risk (see electronic supplementary material S5) may limit the detection of within-individual effects, and so we encourage research aimed at further disentangling plastic effects of predation risk on offspring or adults, respectively. Our results also highlight a predominance of risk-avoiding females (longer FIDs) on islands with high nest predation risk. We hypothesize that the observed non-random distribution of risk-taking phenotypes may be linked with habitat-specific directional selection. Open treeless islands have previously been shown to suffer from higher nest predation risk [34], and to offer less concealed nest-sites, in which females display longer FIDs ([93]; this study). Importantly, in open nest-sites, higher stress-sensitivity (commonly associated with risk-avoidance) has been shown to be associated with lower nest depredation under high predation threat [56]. Consequently, one may expect reproductive selection for risk-

avoiding phenotypes on sparsely vegetated islands characterized by high nest predation risk. As the low repeatability of island-specific adult predation risk ($r=0.103$, electronic supplementary material S6) may render it difficult to detect a significant among-individual effect, we encourage future studies using more replicates to confirm the absence of an among-individual effect of predation risk on adults on FID.

5. Conclusion

Our long-term study showed that risk-taking propensity can increase at the population level despite increasing predation pressure, likely due to compensatory changes in nest-site selection and breeder quality overcoming the direct effects of predation threat on prey behaviour. Nevertheless, at the individual level, female eiders did show the expected increase in FID with increasing predation risk. Importantly, our results suggest that the underlying mechanism—behavioural plasticity or selective (dis)appearance—may differ depending on the target of predation (adults or offspring), in line with the theoretically predicted reluctance of parents in long-lived species to trade their own survival for that of their offspring. In addition, greater propensity for risk-taking in concealed nests was likely a consequence of personality-matching [95], rather than mirroring individual plasticity. We also highlighted that older and better-condition females engaged in greater risk-taking. An important insight arises from the current backdrop of unprecedented increase in predation risk. Selective (dis)appearance effects are typically underestimated in captive populations or in species experiencing low-extrinsic mortality [35]. By contrast, our results suggest that individual quality-dependent mortality and reproductive output may be important drivers of individual plasticity in anti-predator responses in natural populations experiencing high predation risk and the strength of predator-induced selection may even

affect the direction of selection [99]. Important next steps will be to elucidate the lifetime fitness consequences of differential risk-taking strategies under variable predation risk and, as highlighted by our study, such endeavours should strive to isolate the effects of adult and nest predation in wild animals.

Ethics. Handling of animals was reviewed and approved by the National Animal Experiment Board [permit number ESAVI/1697/04.10.03/2012] and complied with the regulations of Tvärminne Zoological Station.

Data accessibility. The data supporting this study are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.m37pvm59> [100].

Electronic supplementary material is available online [101].

Authors' contributions. B.M.: conceptualization, data curation, formal analysis, investigation, visualization, writing—original draft and writing—review and editing; F.A.: conceptualization, investigation, supervision and writing—review and editing; K.J.: conceptualization, funding acquisition, investigation and writing—review and editing; B.S.: investigation and writing—review and editing; E.L.: data curation and investigation; M.Ö.: conceptualization, funding acquisition, investigation, project administration, supervision and writing—review and editing.

All authors gave final approval for publication and agreed to be held accountable for the work performed therein.

Conflict of interest declaration. We declare we have no competing interests.

Funding. This research was supported by the Swedish Cultural Foundation in Finland (grant nos. 17/3317, 16/1476, 15/3296, 14/2657, 13/2654, 138139, 149014, 158026, 168333 and 177733 to M.Ö.) and the Academy of Finland (grant nos. 266208 to K.J. and 128039 to M.Ö.). K.J. was funded by grants from Sophie von Julins Stiftelse and from the Ministry of Agriculture and Forestry of Finland (grant to the Nature and Game Management Trust Finland).

Acknowledgements. We thank Tvärminne Zoological Station for providing facilities. We thank all field assistants who helped collecting the data. We thank Coraline Bichet for fruitful discussions on within-subject centring. Handling of animals was reviewed and approved by the National Animal Experiment Board [permit number ESAVI/1697/04.10.03/2012] and complied with the regulations of Tvärminne Zoological Station.

References

- Williams GC. 1966 Natural selection, the costs of reproduction, and a refinement of Lack's principle. *Am. Nat.* **100**, 687–690. (doi:10.1086/282461)
- Reznick D. 1985 Costs of reproduction: an evaluation of the empirical evidence. *Oikos* **44**, 257–267. (doi:10.2307/3544698)
- Sibly RM, Witt CC, Wright NA, Venditti C, Jetz W, Brown JH. 2012 Energetics, lifestyle, and reproduction in birds. *Proc. Natl Acad. Sci. USA* **109**, 10 937–10 941. (doi:10.1073/pnas.1206512109)
- Magnhagen C. 1991 Predation risk as a cost of reproduction. *Trends Ecol. Evol.* **6**, 183–186. (doi:10.1016/0169-5347(91)90210-0)
- Sih A. 1994 Predation risk and the evolutionary ecology of reproductive behaviour. *J. Fish Biol.* **45**, 111–130. (doi:10.1111/j.1095-8649.1994.tb01087.x)
- Lima SL. 1998 Nonlethal effects in the ecology of predator-prey interactions. *BioScience* **48**, 25–34. (doi:10.2307/1313225)
- Clark CW. 1994 Antipredator behavior and the asset-protection principle. *Behav. Ecol.* **5**, 159–170. (doi:10.1093/beheco/5.2.159)
- Jolly CJ, Webb JK, Phillips BL. 2018 The perils of paradise: an endangered species conserved on an island loses antipredator behaviours within 13 generations. *Biol. Lett.* **14**, 20180222. (doi:10.1098/rsbl.2018.0222)
- Biro PA, Stamps JA. 2008 Are animal personality traits linked to life-history productivity? *Trends Ecol. Evol.* **23**, 361–368. (doi:10.1016/j.tree.2008.04.003)
- Hamel S, Gaillard JM, Festa-Bianchet M, Côté SD. 2009 Individual quality, early-life conditions, and reproductive success in contrasted populations of large herbivores. *Ecology* **90**, 1981–1995. (doi:10.1890/08-0596.1)
- Genovart M, Negre N, Tavecchia G, Bistuer A, Parnal L, Oro D. 2010 The young, the weak and the sick: evidence of natural selection by predation. *PLoS ONE* **5**, e9774. (doi:10.1371/journal.pone.0009774)
- Biro PA, Abrahams MV, Post JR, Parkinson EA. 2004 Predators select against high growth rates and risk-taking behaviour in domestic trout populations. *Proc. R. Soc. Lond. B* **271**, 2233–2237. (doi:10.1098/rspb.2004.2861)
- Dingemanse NJ, Both C, Drent PJ, Tinbergen JM. 2004 Fitness consequences of avian personalities in a fluctuating environment. *Proc. R. Soc. Lond. B* **271**, 847–852. (doi:10.1098/rspb.2004.2680)
- Smith BR, Blumstein DT. 2008 Fitness consequences of personality: a meta-analysis. *Behav. Ecol.* **19**, 448–455. (doi:10.1093/beheco/arm144)
- Niemela PT, Lattenkamp EZ, Dingemanse NJ. 2015 Personality-related survival and sampling bias in wild cricket nymphs. *Behav. Ecol.* **26**, 936–946. (doi:10.1093/beheco/arv036)
- Ghalambor CK, Martin TE. 2001 Fecundity-survival trade-offs and parental risk-taking in birds. *Science* **292**, 494–497. (doi:10.1126/science.1059379)
- Fontaine JJ, Martin TE. 2006 Parent birds assess nest predation risk and adjust their reproductive strategies. *Ecol. Lett.* **9**, 428–434. (doi:10.1111/j.1461-0248.2006.00892.x)
- van de Pol M, Verhulst S. 2006 Age-dependent traits: a new statistical model to separate within- and between-individual effects. *Am. Nat.* **167**, 766–773. (doi:10.1086/503331)

19. Blumstein DT. 2019 What chasing birds can teach us about predation risk effects: past insights and future directions. *J. Ornithol.* **160**, 587–592. (doi:10.1007/s10336-019-01634-1)
20. Ydenberg RC, Dill LM. 1986 The economics of fleeing from predators. In *Advances in the study of behavior*, pp. 229–249. Amsterdam, The Netherlands: Elsevier.
21. Blumstein DT, Samia DS, Cooper WE. 2016 Escape behavior: dynamic decisions and a growing consensus. *Curr. Opin. Behav. Sci.* **12**, 24–29. (doi:10.1016/j.cobeha.2016.08.006)
22. Carrete M, Tella JL. 2010 Individual consistency in flight initiation distances in burrowing owls: a new hypothesis on disturbance-induced habitat selection. *Biol. Lett.* **6**, 167–170. (doi:10.1098/rsbl.2009.0739)
23. Møller AP, Garamszegi LZ. 2012 Between individual variation in risk-taking behavior and its life history consequences. *Behav. Ecol.* **23**, 843–853. (doi:10.1093/beheco/ars040)
24. Cabrera D, Andres D, McLoughlin PD, Debeffe L, Medill SA, Wilson AJ, Poissant J. 2017 Island tameness and the repeatability of flight initiation distance in a large herbivore. *Can. J. Zool.* **95**, 771–778. (doi:10.1139/cjz-2016-0305)
25. Strong JS, Weladji RB, Holand Ø, Røed KH, Nieminen M. 2017 Personality and fitness consequences of flight initiation distance and mating behavior in subdominant male reindeer (*Rangifer tarandus*). *Ethology* **123**, 484–492. (doi:10.1111/eth.12619)
26. de Jong ME, Nicolaus M, Fokkema RW, Loonen MJJE. 2021 State dependence explains individual variation in nest defence behaviour in a long-lived bird. *J. Anim. Ecol.* **90**, 809–819. (doi:10.1111/1365-2656.13411)
27. Broom M, Ruxton GD. 2005 You can run—or you can hide: optimal strategies for cryptic prey against pursuit predators. *Behav. Ecol.* **16**, 534–540. (doi:10.1093/beheco/ari024)
28. Albrecht T, Klvaňa P. 2004 Nest crypsis, reproductive value of a clutch and escape decisions in incubating female mallards *Anas platyrhynchos*. *Ethology* **110**, 603–613. (doi:10.1111/j.1439-0310.2004.00992.x)
29. Wolf M, van Doorn GS, Leimar O, Weissing FJ. 2007 Life-history trade-offs favour the evolution of animal personalities. *Nature* **447**, 581–584. (doi:10.1038/nature05835)
30. Dowling L, Bonier F. 2018 Should I stay, or should I go: modeling optimal flight initiation distance in nesting birds. *PLoS ONE* **13**, e0208210. (doi:10.1371/journal.pone.0208210)
31. Piersma T, Drent J. 2003 Phenotypic flexibility and the evolution of organismal design. *Trends Ecol. Evol.* **18**, 228–233. (doi:10.1016/S0169-5347(03)00036-3)
32. Edelaar P, Siepielski AM, Clobert J. 2008 Matching habitat choice causes directed gene flow: a neglected dimension in ecology and evolution. *Evolution* **62**, 2462–2472. (doi:10.1111/j.1558-5646.2008.00459.x)
33. Abbey-Lee RN, Mathot KJ, Dingemanse NJ. 2016 Behavioral and morphological responses to perceived predation risk: a field experiment in passerines. *Behav. Ecol.* **27**, 857–864. (doi:10.1093/beheco/aru228)
34. Öst M, Lindén A, Karell P, Ramula S, Kilpi M. 2018 To breed or not to breed: drivers of intermittent breeding in a seabird under increasing predation risk and male bias. *Oecologia* **188**, 129–138. (doi:10.1007/s00442-018-4176-5)
35. Hämäläinen A, Dammhahn M, Aujard F, Eberle M, Hardy I, Kappeler PM, Perret M, Schliehe-Diecks S, Kraus C. 2014 Senescence or selective disappearance? Age trajectories of body mass in wild and captive populations of a small-bodied primate. *Proc. R. Soc. B* **281**, 20140830. (doi:10.1098/rspb.2014.0830)
36. Korschgen CE. 1977 Breeding stress of female eiders in Maine. *J. Wildl. Manage.* **41**, 360–373. (doi:10.2307/3800505)
37. Bolduc F, Guillemette M. 2003 Human disturbance and nesting success of common eiders: interaction between visitors and gulls. *Biol. Conserv.* **110**, 77–83. (doi:10.1016/S0006-3207(02)00178-7)
38. Kilpi M, Lindström K. 1997 Habitat-specific clutch size and cost of incubation in common eiders, *Somateria mollissima*. *Oecologia* **111**, 297–301. (doi:10.1007/s004420050238)
39. Hanssen SA, Erikstad KE, Johnsen V, Bustnes JO. 2003 Differential investment and costs during avian incubation determined by individual quality: an experimental study of the common eider (*Somateria mollissima*). *Proc. R. Soc. Lond. B* **270**, 531–537. (doi:10.1098/rspb.2002.2262)
40. Jaatinen K, Seltmann MW, Hollmén T, Atkinson S, Mashburn K, Öst M. 2013 Context dependency of baseline glucocorticoids as indicators of individual quality in a capital breeder. *Gen. Comp. Endocrinol.* **191**, 231–238. (doi:10.1016/j.ygcen.2013.06.022)
41. Coulson J. 1984 The population dynamics of the eider duck *Somateria mollissima* and evidence of extensive non-breeding by adult ducks. *Ibis* **126**, 525–543. (doi:10.1111/j.1474-919X.1984.tb02078.x)
42. Öst M, Lehikoinen A, Jaatinen K, Kilpi M. 2011 Causes and consequences of fine-scale breeding dispersal in a female-philopatric species. *Oecologia* **166**, 327–336. (doi:10.1007/s00442-010-1855-2)
43. Coulson JC. 2016 A review of philopatry in seabirds and comparisons with other waterbird species. *Waterbirds* **39**, 229–240. (doi:10.1675/063.039.0302)
44. Nicol-Harper A, Wood KA, Diamond AW, Major HL, Petersen A, Tertitski G, Doncaster CP, Ezard THG, Hilton GM. 2021 Vital rate estimates for the common eider *Somateria mollissima*, a data-rich exemplar of the seaduck tribe. *Ecol. Solutions Evid.* **2**, e12108. (doi:10.1002/2688-8319.12108)
45. Hario M, Rintala J. 2009 Age of first breeding in the common eider *Somateria m. mollissima* population in the northern Baltic Sea. *Ornis Fennica* **86**, 81–88.
46. Jaatinen K, Öst M. 2011 Experience attracts: the role of age in the formation of cooperative brood-rearing coalitions in eiders. *Anim. Behav.* **81**, 1289–1294. (doi:10.1016/j.anbehav.2011.03.020)
47. Criscuolo F, Gauthier-Clerc M, Gabrielsen GW, Le Maho Y. 2000 Recess behaviour of the incubating common eider *Somateria mollissima*. *Polar Biol.* **23**, 571–574. (doi:10.1007/s003000000123)
48. Öst M, Steele BB. 2010 Age-specific nest-site preference and success in eiders. *Oecologia* **162**, 59–69. (doi:10.1007/s00442-009-1444-4)
49. Frid A, Dill L. 2002 Human-caused disturbance stimuli as a form of predation risk. *Conserv. Ecol.* **6**, 11. (doi:10.5751/ES-00404-060111)
50. Seltmann MW, Öst M, Jaatinen K, Atkinson S, Mashburn K, Hollmén T. 2012 Stress responsiveness, age and body condition interactively affect flight initiation distance in breeding female eiders. *Anim. Behav.* **84**, 889–896. (doi:10.1016/j.anbehav.2012.07.012)
51. Forbes MRL, Clark RG, Weatherhead PJ, Armstrong T. 1994 Risk-taking by female ducks: intra- and interspecific tests of nest defense theory. *Behav. Ecol. Sociobiol.* **34**, 79–85. (doi:10.1007/BF00164178)
52. Osiejuk TS, Kuczyński L. 2007 Factors affecting flushing distance in incubating female greylag geese *Anser anser*. *Wildl. Biol.* **13**, 11–18. (doi:10.2981/0909-6396(2007)13[11:FAFDII]2.0.CO;2)
53. Götmark F, Åhlund M. 1984 Do field observers attract nest predators and influence nesting success of common eiders? *J. Wildl. Manage.* **48**, 381. (doi:10.2307/3801169)
54. Abràmoff MD, Magalhães PJ, Ram SJ. 2004 Image processing with ImageJ. *Biophotonics Int.* **11**, 36–42.
55. Öst M, Lehikoinen A, Jaatinen K. 2022 Top–down effects override climate forcing on reproductive success in a declining sea duck. *Oikos* **2022**, e08762. (doi:10.1111/oik.08762)
56. Jaatinen K, Seltmann MW, Öst M. 2014 Context-dependent stress responses and their connections to fitness in a landscape of fear. *J. Zool.* **294**, 147–153. (doi:10.1111/jzo.12169)
57. Mohring B, Angelier F, Jaatinen K, Parenteau C, Öst M. 2021 Parental investment under predation threat in incubating common eiders (*Somateria mollissima*): a hormonal perspective. *Front. Ecol. Evol.* **9**, 637561. (doi:10.3389/fevo.2021.637561)
58. Öst M, Bäck A. 2003 Spatial structure and parental aggression in eider broods. *Anim. Behav.* **66**, 1069–1075. (doi:10.1006/anbe.2003.2300)
59. R Core Team. 2020 *R: a language and environment for statistical computing*. Austria, Vienna: R Foundation for Statistical Computing.
60. Lüdtke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. 2021 performance: An R package for assessment, comparison and testing of statistical models. *JOSS* **6**, 3139. (doi:10.21105/joss.03139)
61. Allison P. 2012 When can you safely ignore multicollinearity. *Stat. Horizons* **5**, 1–2.
62. van de Pol M, Wright J. 2009 A simple method for distinguishing within- versus between-subject effects using mixed models. *Anim. Behav.* **77**, 753–758. (doi:10.1016/j.anbehav.2008.11.006)
63. Stoffel M, Nakagawa S, Schielzeth H, Goslee S. 2017 rptR: repeatability estimation and variance decomposition by generalized linear mixed-effects

- models. *Methods Ecol. Evol.* **8**, 1639–1644. (doi:10.1111/2041-210X.12797)
64. Salomons HM, Mulder GA, van de Zande L, Haussmann MF, Linskens MHK, Verhulst S. 2009 Telomere shortening and survival in free-living corvids. *Proc. R. Soc. B* **276**, 3157–3165. (doi:10.1098/rspb.2009.0517)
65. Berger S, Wikelski M, Romero LM, Kalko EKV, Rödl T. 2007 Behavioral and physiological adjustments to new predators in an endemic island species, the Galápagos marine iguana. *Horm. Behav.* **52**, 653–663. (doi:10.1016/j.yhbeh.2007.08.004)
66. Cooper Jr WE, Hawlena D, Pérez-Mellado V. 2009 Effects of predation risk factors on escape behavior by Balearic lizards (*Podarcis lilfordi*) in relation to optimal escape theory. *Amphib. Reptilia* **30**, 99–110. (doi:10.1163/156853809787392720)
67. Zamora-Camacho FJ, García-Astilleros J, Aragón P. 2018 Does predation risk outweigh the costs of lost feeding opportunities or does it generate a behavioural trade-off? A case study with Iberian ribbed newt larvae. *Biol. J. Linn. Soc. Lond.* **125**, 741–749. (doi:10.1093/biolinnean/bly155)
68. Dingemanse NJ, Dochtermann NA. 2013 Quantifying individual variation in behaviour: mixed-effect modelling approaches. *J. Anim. Ecol.* **82**, 39–54. (doi:10.1111/1365-2656.12013)
69. Kroos J, Öst M, Karell P, Jaatinen K, Kilpi M. 2012 Philopatric predisposition to predation-induced ecological traps: habitat-dependent mortality of breeding eiders. *Oecologia* **170**, 979–986. (doi:10.1007/s00442-012-2378-9)
70. Lima SL. 2009 Predators and the breeding bird: behavioral and reproductive flexibility under the risk of predation. *Biol. Rev.* **84**, 485–513. (doi:10.1111/j.1469-185X.2009.00085.x)
71. Dulude-de Broin F, Hamel S, Mastromonaco GF, Côté SD. 2020 Predation risk and mountain goat reproduction: evidence for stress-induced breeding suppression in a wild ungulate. *Funct. Ecol.* **34**, 1003–1014. (doi:10.1111/1365-2435.13514)
72. Chastel O, Weimerskirch H, Jouventin P. 1995 Body condition and seabird reproductive performance: a study of three petrel species. *Ecology* **76**, 2240–2246. (doi:10.2307/1941698)
73. Wendeln H, Becker PH. 1999 Effects of parental quality and effort on the reproduction of common terns. *J. Anim. Ecol.* **68**, 205–214. (doi:10.1046/j.1365-2656.1999.00276.x)
74. Lehikoinen A, Jaatinen K, Öst M. 2010 Do female ornaments indicate quality in eider ducks? *Biol. Lett.* **6**, 225–228. (doi:10.1098/rsbl.2009.0744)
75. Bell AM, Hankison SJ, Laskowski KL. 2009 The repeatability of behaviour: a meta-analysis. *Anim. Behav.* **77**, 771–783. (doi:10.1016/j.anbehav.2008.12.022)
76. Kenward R. 1978 Hawks and doves: factors affecting success and selection in goshawk attacks on woodpigeons. *J. Anim. Ecol.* **47**, 449–460. (doi:10.2307/3793)
77. Martín J, de Neve L, Polo V, Fargallo JA, Soler M. 2006 Health-dependent vulnerability to predation affects escape responses of unguarded chinstrap penguin chicks. *Behav. Ecol. Sociobiol.* **60**, 778–784. (doi:10.1007/s00265-006-0221-1)
78. Luttbeg B, Sih A. 2010 Risk, resources and state-dependent adaptive behavioural syndromes. *Phil. Trans. R. Soc. B* **365**, 3977–3990. (doi:10.1098/rstb.2010.0207)
79. Tablado Z, Jenni L. 2017 Determinants of uncertainty in wildlife responses to human disturbance: modulators of wildlife response to recreation. *Biol. Rev.* **92**, 216–233. (doi:10.1111/brv.12224)
80. Curio E. 1983 Why do young birds reproduce less well? *Ibis* **125**, 400–404. (doi:10.1111/j.1474-919X.1983.tb03130.x)
81. Candolin U. 1998 Reproduction under predation risk and the trade-off between current and future reproduction in the threespine stickleback. *Proc. R. Soc. Lond. B* **265**, 1171–1175. (doi:10.1098/rspb.1998.0415)
82. Daan S, Dijkstra C, Drent R, Meijer T. 1988 Food supply and the annual timing of avian reproduction. In *Proc. of the International Ornithological Congress*, pp. 392–407. Ottawa, Canada: University of Ottawa Press Ottawa.
83. Erikstad KE, Bustnes JO, Moum T. 1993 Clutch-size determination in precocial birds: a study of the common eider. *The Auk* **110**, 623–628. (doi:10.2307/4088426)
84. Gatti RC. 1983 Incubation weight loss in the mallard. *Can. J. Zool.* **61**, 565–569. (doi:10.1139/z83-076)
85. Garbus SE *et al.* 2018 Incubation behaviour of common eiders *Somateria mollissima* in the Central Baltic: nest attendance and loss in body mass. *Acrocephalus* **39**, 91–100. (doi:10.1515/acro-2018-0008)
86. Öst M, Clark CW, Kilpi M, Ydenberg R. 2007 Parental effort and reproductive skew in coalitions of brood rearing female common eiders. *Am. Nat.* **169**, 73–86. (doi:10.1086/510213)
87. Waldeck P, Kilpi M, Andersson M, Öst M. 2004 Brood parasitism in a population of common eider (*Somateria mollissima*). *Behaviour* **141**, 725–739. (doi:10.1163/1568539042245132)
88. Öst M, Wickman M, Matulionis E, Steele B. 2008 Habitat-specific clutch size and cost of incubation in eiders reconsidered. *Oecologia* **158**, 205–216. (doi:10.1007/s00442-008-1139-2)
89. Altmann M. 1958 Social integration of the moose calf. *Anim. Behav.* **6**, 155–159. (doi:10.1016/0003-3472(58)90045-9)
90. Stankowich T, Blumstein DT. 2005 Fear in animals: a meta-analysis and review of risk assessment. *Proc. R. Soc. B* **272**, 2627–2634. (doi:10.1098/rspb.2005.3251)
91. Stankowich T. 2008 Ungulate flight responses to human disturbance: a review and meta-analysis. *Biol. Conserv.* **141**, 2159–2173. (doi:10.1016/j.biocon.2008.06.026)
92. Miller V, Abraham KF, Nol E. 2013 Factors affecting the responses of female Canada geese to disturbance during incubation. *J. Field Ornithol.* **84**, 171–180. (doi:10.1111/jof.12015)
93. Seltmann MW, Jaatinen K, Steele BB, Öst M. 2014 Boldness and stress responsiveness as drivers of nest-site selection in a ground-nesting bird. *Ethology* **120**, 77–89. (doi:10.1111/eth.12181)
94. Wilson-Aggarwal JK, Troscianko JT, Stevens M, Spottiswoode CN. 2016 Escape distance in ground-nesting birds differs with individual level of camouflage. *Am. Nat.* **188**, 231–239. (doi:10.1086/687254)
95. Holtmann B, Santos ESA, Lara CE, Nakagawa S. 2017 Personality-matching habitat choice, rather than behavioural plasticity, is a likely driver of a phenotype–environment covariance. *Proc. R. Soc. B* **284**, 20170943. (doi:10.1098/rspb.2017.0943)
96. D'Alba L, Spencer KA, Nager RG, Monaghan P. 2011 State dependent effects of elevated hormone: nest site quality, corticosterone levels and reproductive performance in the common eider. *Gen. Comp. Endocrinol.* **172**, 218–224. (doi:10.1016/j.ygcen.2011.03.006)
97. Gaillard JM, Yoccoz NG. 2003 Temporal variation in survival of mammals: a case of environmental canalization? *Ecology* **84**, 3294–3306. (doi:10.1890/02-0409)
98. Hamel S, Gaillard JM, Yoccoz NG, Loison A, Bonenfant C, Descamps S. 2010 Fitness costs of reproduction depend on life speed. *Ecol. Lett.* **13**, 915–935. (doi:10.1111/j.1461-0248.2010.01478.x)
99. Jaatinen K, Möller AP, Öst M. 2019 Annual variation in predation risk is related to the direction of selection for brain size in the wild. *Sci. Rep.* **9**, 11847. (doi:10.1038/s41598-019-48153-w)
100. Mohring B, Angelier F, Jaatinen K, Steele B, Lönnerberg E, Öst M. 2022 Data from: Drivers of within- and among-individual variation in risk-taking behaviour during reproduction in a long-lived bird. Dryad Digital Repository. (doi:10.5061/dryad.m37pvm59)
101. Mohring B, Angelier F, Jaatinen K, Steele B, Lönnerberg E, Öst M. 2022 Drivers of within- and among-individual variation in risk-taking behaviour during reproduction in a long-lived bird. Figshare. (doi:10.6084/m9.figshare.c.6182022)